

Supplementary Table 2. Subgroup analyses in the peripheral vertigo group

Characteristic	AUVP/VN (n=17)	BPPV (n=21)	MD (n=7)	p-value
Age (yr)	62.35±14.54	66.48±12.69	61.71±11.76	0.56
Male sex	9 (52.9)	7 (33.3)	3 (42.9)	0.48
Pulse wave velocity				
Right	11.53±2.87	12.38±2.81	13.44±2.57	0.41
Left	11.37±3.39	13.46±3.60	13.56±2.19	0.20
Mean	11.45±3.08	12.92±3.02	13.50±2.25	0.28
Ankle-brachial index				
Right	1.18±0.14	1.19±0.10	1.09±0.12	0.18
Left	1.14±0.12	1.15±0.08	1.18±0.11	0.74
Mean	1.16±0.12	1.17±0.07	1.14±0.10	0.70
Hypertension	8 (47.1)	6 (28.6)	3 (42.9)	0.48
Diabetes mellitus	3 (17.6)	3 (14.3)	0 (0)	0.50
Atrial fibrillation	1 (6.3)	1 (4.8)	2 (28.6)	0.15
Previous stroke	0 (0)	1 (4.8)	0 (0)	0.56
Dyslipidemia	4 (23.5)	4 (19.0)	2 (28.6)	0.86
Smoking	5 (29.4)	1 (4.8)	1 (14.3)	0.15

Values are presented as mean±standard deviation or number (%).

AUVP, acute unilateral vestibulopathy; VN, vestibular neuritis; BPPV, benign paroxysmal positional vertigo; MD, Ménière's disease.