

**Supplementary Table 3.** Subgroup analyses in the central vertigo group

Characteristic	AICA (n=21)	PICA (n=11)	SCA (n=16)	p-value
Age (yr)	63.71±10.92	57.00±13.41	68.00±12.78	0.08
Male sex	17 (81.1)	7 (63.6)	7 (43.8)	0.06
Pulse wave velocity				
Right	16.76±4.77	18.88±4.81	17.00±5.33	0.57
Left	18.63±5.28	18.28±6.54	19.01±6.20	0.95
Mean	17.75±4.31	17.95±5.97	18.92±6.93	0.88
Ankle-brachial index				
Right	1.16±0.08	1.13±0.22	1.10±0.14	0.45
Left	1.14±0.10	1.16±0.10	1.13±0.11	0.73
Mean	1.15±0.08	1.14±0.14	1.11±0.11	0.57
Hypertension	11 (52.4)	6 (54.5)	9 (56.2)	0.97
Diabetes mellitus	7 (33.3)	3 (27.3)	7 (43.8)	0.66
Atrial fibrillation	1 (4.8)	1 (9.1)	3 (18.8)	0.38
Previous stroke	3 (14.3)	1 (9.1)	1 (6.2)	0.72
Dyslipidemia	10 (47.6)	7 (63.6)	10 (62.5)	0.24
Smoking	9 (42.9)	4 (36.4)	5 (31.3)	0.69

AICA, anterior cerebellar arterial infarction; PICA, posterior cerebellar arterial infarction.